

Hindawi Publishing Corporation
Journal of Inequalities and Applications
Volume 2010, Article ID 546015, 4 pages
doi:10.1155/2010/546015

Research Article

Alon-Babai-Suzuki's Conjecture Related to Binary Codes in Nonmodular Version

K.-W. Hwang,¹ T. Kim,² L. C. Jang,³ P. Kim,⁴ and Gyoyong Sohn⁵

¹ Department of Mathematics, Donga-A University, Pusan 604-714, South Korea

² Division of General Edu.-Math., Kwangwoon University, Seoul 139-701, South Korea

³ Department of Mathematics and Computer Science, Konkook University, Chungju 139-701, South Korea

⁴ Department of Mathematics, Kyungpook National University, Taegu 702-701, South Korea

⁵ Department of Computer Science, Chungbuk National University, Cheongju 361-763, South Korea

Correspondence should be addressed to K.-W. Hwang, khwang7@kookmin.ac.kr

Received 23 August 2009; Accepted 22 January 2010

Academic Editor: Ram N. Mohapatra

Copyright © 2010 K.-W. Hwang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Let $K = \{k_1, k_2, \dots, k_r\}$ and $L = \{l_1, l_2, \dots, l_s\}$ be sets of nonnegative integers. Let $\mathcal{F} = \{F_1, F_2, \dots, F_m\}$ be a family of subsets of $[n]$ with $|F_i| \in K$ for each i and $|F_i \cap F_j| \in L$ for any $i \neq j$. Every subset F_e of $[n]$ can be represented by a binary code $\mathbf{a} = (a_1, a_2, \dots, a_n)$ such that $a_i = 1$ if $i \in F_e$ and $a_i = 0$ if $i \notin F_e$. Alon et al. made a conjecture in 1991 in modular version. We prove Alon-Babai-Sukuki's Conjecture in nonmodular version. For any K and L with $n \geq s + \max k_i$, $|\mathcal{F}| \leq \binom{n-1}{s} + \binom{n-1}{s-1} + \dots + \binom{n-1}{s-2r+1}$.

1. Introduction

In this paper, \mathcal{F} stands for a family of subsets of $[n] = \{1, 2, \dots, n\}$, $K = \{k_1, \dots, k_r\}$, and $L = \{l_1, \dots, l_s\}$, where $|F_i| \in K$ for all $F_i \in \mathcal{F}$, $|F_i \cap F_j| \in L$ for all $F_i, F_j \in \mathcal{F}, i \neq j$. The variable x will stand as a shorthand for the n -dimensional vector variable (x_1, x_2, \dots, x_n) . Also, since these variables will take the values only 0 and 1, all the polynomials we will work with will be reduced modulo the relation $x_i^2 = x_i$. We define the characteristic vector $v_i = (v_{i1}, v_{i2}, \dots, v_{in})$ of F_i such that $v_{ij} = 1$ if $j \in F_i$ and $v_{ij} = 0$ if $j \notin F_i$. We will present some results in this paper that give upper bounds on the size of \mathcal{F} under various conditions. Below is a list of related results by others.

Theorem 1.1 (Ray-Chaudhuri and Wilson [1]). *If $K = \{k\}$, and L is any set of nonnegative integers with $k > \max l_j$, then $|\mathcal{F}| \leq \binom{n}{s}$.*

Theorem 1.2 (Alon et al. [2]). *If K and L are two sets of nonnegative integers with $k_i > s - r$, for every i , then $|\mathcal{F}| \leq \binom{n}{s} + \binom{n}{s-1} + \cdots + \binom{n}{s-r+1}$.*

Theorem 1.3 (Snevily [3]). *If K and L are any sets such that $\min k_i > \max l_j$, then $|\mathcal{F}| \leq \binom{n-1}{s} + \binom{n-1}{s-1} + \cdots + \binom{n-1}{0}$.*

Theorem 1.4 (Snevily [4]). *Let K and L be sets of nonnegative integers such that $\min k_i > \max l_j$. Then, $|\mathcal{F}| \leq \binom{n-1}{s} + \binom{n-1}{s-1} + \cdots + \binom{n-1}{s-2r+1}$.*

Conjecture 1.5 (Snevily [5]). *For any K and L with $\min k_i > \max l_j$, $|\mathcal{F}| \leq \binom{n}{s}$.*

In the same paper in which he stated the above conjecture, Snevily mentions that it seems hard to prove the above bound and states the following weaker conjecture.

Conjecture 1.6 (Snevily [5]). *For any K and L with $\min k_i > \max l_j$, $|\mathcal{F}| \leq \binom{n-1}{s} + \binom{n-1}{s-1} + \cdots + \binom{n-1}{s-r}$.*

Hwang and Sheikh [6] proved the bound of Conjecture 1.6 when K is a consecutive set. The second theorem we prove is a special case of Conjecture 1.6 with the extra condition that $\bigcap_{i=1}^m F_i \neq \emptyset$. These two theorems are stated hereunder.

Theorem 1.7 (Hwang and Sheikh [6]). *Let $K = \{k_1, k_2, \dots, k_r\}$ where $k_i = k_1 + i - 1$, $k_1 > s - r$, and $L = \{l_1, l_2, \dots, l_s\}$. Let $\mathcal{F} = \{F_1, F_2, \dots, F_m\}$ be such that $|F_i| \in K$ for each i , $|F_i| \notin L$, and $|F_i \cap F_j| \in L$ for any $i \neq j$. Then $|\mathcal{F}| \leq \binom{n-1}{s} + \binom{n-1}{s-1} + \cdots + \binom{n-1}{s-r}$.*

Theorem 1.8 (Hwang and Sheikh [6]). *Let $K = \{k_1, k_2, \dots, k_r\}$, $L = \{l_1, l_2, \dots, l_s\}$, and $\mathcal{F} = \{F_1, F_2, \dots, F_m\}$ be such that $|F_i| \in K$ for each i , $|F_i \cap F_j| \in L$ for any $i \neq j$, and $k_i > s - r$. If $\bigcap_{i=1}^m F_i \neq \emptyset$, then $|\mathcal{F}| \leq \binom{n-1}{s} + \binom{n-1}{s-1} + \cdots + \binom{n-1}{s-r}$.*

Theorem 1.9 (Alon et al. [2]). *Let K and L be subsets of $\{0, 1, \dots, p-1\}$ such that $K \cap L = \emptyset$, where p is a prime and $\mathcal{F} = \{F_1, F_2, \dots, F_m\}$ a family of subsets of $[n]$ such that $|F_i| \pmod{p} \in K$ for all $F_i \in \mathcal{F}$ and $|F_i \cap F_j| \pmod{p} \in L$ for $i \neq j$. If $r(s-r+1) \leq p-1$, and $n \geq s + \max k_i$, then $|\mathcal{F}| \leq \binom{n}{s} + \binom{n}{s-1} + \cdots + \binom{n}{s-r+1}$.*

Conjecture 1.10 (Alon et al. [2]). *Let K and L be subsets of $\{0, 1, \dots, p-1\}$ such that $K \cap L = \emptyset$, where p is a prime and $\mathcal{F} = \{F_1, F_2, \dots, F_m\}$ a family of subsets of $[n]$ such that $|F_i| \pmod{p} \in K$ for all $F_i \in \mathcal{F}$ and $|F_i \cap F_j| \pmod{p} \in L$ for $i \neq j$. If $n \geq s + \max k_i$, then $|\mathcal{F}| \leq \binom{n}{s} + \binom{n}{s-1} + \cdots + \binom{n}{s-r+1}$.*

In [2], Alon et al. proved their conjectured bound under the extra conditions that $r(s-r+1) \leq p-1$ and $n \geq s + \max k_i$. Qian and Ray-Chaudhuri [7] proved that if $n > 2s-r$ instead of $n \geq s + \max k_i$, then the above bound holds.

We prove an Alon-Babai-Suzuki's conjecture in non-modular version.

Theorem 1.11. *Let $K = \{k_1, k_2, \dots, k_r\}$, $L = \{l_1, l_2, \dots, l_s\}$ be two sets of nonnegative integers and let $\mathcal{F} = \{F_1, F_2, \dots, F_m\}$ be such that $|F_i| \in K$ for each i , $|F_i \cap F_j| \in L$ for any $i \neq j$, and $n \geq s + \max_i |F_i|$. then $|\mathcal{F}| \leq \binom{n}{s} + \binom{n}{s-1} + \cdots + \binom{n}{s-r+1}$.*

2. Proof of Theorem

Proof of Theorem 1.11. For each $F_i \in \mathcal{F}$, consider the polynomial

$$f_i(x) = \prod_{\substack{j \\ l_j < |F_i|}} (v_i \cdot x - (k_i - l_j)), \quad (2.1)$$

where v_i is the characteristic vector of F_i and v_i^* is the characteristic vector of $F_i^* = F_i - \{1\}$. Let \bar{v}_i be the characteristic vector of F_i^c , and \bar{v}_i^* be the characteristic vector of $(F_i^c)^*$.

We order $\{F_i\}$ by size of F_i , that is, $|F_j| \leq |F_k|$ if $j < k$. We substitute the characteristic vector \bar{v}_i of F_i^c by order of size of F_i . Clearly, $f_i(\bar{v}_i) \neq 0$ for $1 \leq i \leq m$ and $f_i(\bar{v}_j) = 0$ for $1 \leq j < i \leq m$. Assume that

$$\sum_i \alpha_i f_i(x) = 0. \quad (2.2)$$

We prove that $\{f_i(x)\}$ is linearly independent. Assume that this is false. Let i_0 be the smallest index such that $\alpha_{i_0} \neq 0$. We substitute \bar{v}_{i_0} into the above equation. Then we get $\alpha_{i_0} f_{i_0}(\bar{v}_{i_0}) = 0$. We get a contradiction. So $\{f_i(x)\}$ is linearly independent. Let $\mathcal{E} = \{E_1, \dots, E_e\}$ be the family of subsets of $[n]$ with size at most $s - r$, which is ordered by size, that is, $|E_i| \leq |E_j|$ if $i < j$, where $e = \sum_{i=0}^{s-r} \binom{n}{i}$. Let u_i denote the characteristic vector of E_i . We define the multilinear polynomial g_i in n variables for each E_i :

$$g_i(x) = \prod_{l=1}^r \left(\sum_{t=1}^n x_t - (n - k_l) \right) \prod_{j \in E_i} x_j. \quad (2.3)$$

We prove that $\{g_i(x)\}$ is linearly independent. Assume that

$$\sum_i \beta_i g_i(x) = 0. \quad (2.4)$$

Choose the smallest size of E_i . Let u_i be the characteristic vector of E_i . We substitute u_i into the above equation. We know that $g_i(u_i) \neq 0$ and $g_j(u_i) = 0$ for any $i < j$. Since $n \geq s + \max k_i$, we get $\beta_i = 0$. If we follow the same process, then the family $\{g_i(x)\}$ is linearly independent. Next, we prove that $\{f_i(x), g_i(x)\}$ is linearly independent. Now, assume that

$$\sum_i \alpha_i f_i(x) + \sum_i \beta_i g_i(x) = 0. \quad (2.5)$$

Let F_1 be the smallest size of F_i . We substitute the characteristic vector \bar{v}_1 of F_1^c into the above equation. Since $|F_1^c| = n - k_1$, $g_i(\bar{v}_1) = 0$ for all i . We only get $\alpha_1 f_1(\bar{v}_1) = 0$. So $\alpha_1 = 0$. By the same way, choose the smallest size from $\{F_i\}$ after deleting F_1 . We do the same process. We also can get $\alpha_2 = 0$. By the same process, we prove that all $\alpha_i = 0$. We prove that $\{f_i(x), g_i(x)\}$ is linearly independent.

Any polynomial in the set $\{f_i(x), g_i(x)\}$ can be represented by a linear combination of multilinear monomials of degree $\leq s$. The space of such multilinear polynomials has dimension $\sum_{i=0}^s \binom{n}{i}$. We found $|\mathcal{F}| + \sum_{i=0}^{s-r} \binom{n}{i}$ linearly independent polynomials with degree at most s . So $|\mathcal{F}| + \sum_{i=0}^{s-r} \binom{n}{i} \leq \sum_{i=0}^s \binom{n}{i}$. Thus $|\mathcal{F}| \leq \binom{n}{s} + \binom{n}{s-1} + \cdots + \binom{n}{s-r+1}$. \square

Acknowledgments

The authors thank Zoltán Füredi for encouragement to write this paper. The present research has been conducted by the research grant of the Kwangwoon University in 2009.

References

- [1] D. K. Ray-Chaudhuri and R. M. Wilson, "On t -designs," *Osaka Journal of Mathematics*, vol. 12, no. 3, pp. 737–744, 1975.
- [2] N. Alon, L. Babai, and H. Suzuki, "Multilinear polynomials and Frankl-Ray-Chaudhuri-Wilson type intersection theorems," *Journal of Combinatorial Theory. Series A*, vol. 58, no. 2, pp. 165–180, 1991.
- [3] H. S. Snevily, "On generalizations of the de Bruijn-Erdős theorem," *Journal of Combinatorial Theory. Series A*, vol. 68, no. 1, pp. 232–238, 1994.
- [4] H. S. Snevily, "A sharp bound for the number of sets that pairwise intersect at k positive values," *Combinatorica*, vol. 23, no. 3, pp. 527–533, 2003.
- [5] H. S. Snevily, "A generalization of the Ray-Chaudhuri-Wilson theorem," *Journal of Combinatorial Designs*, vol. 3, no. 5, pp. 349–352, 1995.
- [6] K.-W. Hwang and N. Sheikh, "Intersection families and Snevily's conjecture," *European Journal of Combinatorics*, vol. 28, no. 3, pp. 843–847, 2007.
- [7] J. Qian and D. K. Ray-Chaudhuri, "On mod- p Alon-Babai-Suzuki inequality," *Journal of Algebraic Combinatorics*, vol. 12, no. 1, pp. 85–93, 2000.